

# Gravity as a quantum effect on quantum space-time

Harold Steinacker

Department of Physics, University of Vienna

FWF



Humboldt Colleg, EISA, september 20, 2021

(higher-spin) gravity emerges on suitable background within

IKKT = IIB Matrix Model

Ishibashi, Kawai, Kitazawa, Tsuchiya 1996

$$S = \text{Tr}([Y^a, Y^b][Y_a, Y_b] + \bar{\Psi}\Gamma_a[Y^a, \Psi])$$

maximal SUSY, closely related to IIB string theory

- leads to ( $\mathfrak{hs}$ ) gauge theory on suitable quantum space-time, classical action differs from GR
- quantum effects (1-loop)
  - induced Einstein-Hilbert term, gravity \* new! \*
- weak coupling, no holography, no target space compactification (no landscape!)

outline:

- Yang-Mills matrix models & emergent geometry
- fuzzy  $H_n^4$ , covar. space-time  $\mathcal{M}_n^{3,1}$   
linearized fluctuations, no ghosts
- nonlinear regime:  
frame, metric, torsion, covariant eom
- **quantization:** \* new, unpublished \*  
1-loop effective action  
→ Einstein-Hilbert action (+ extras)  
no cosm. const. problem

introductory review: [arXiv:1911.03162](https://arxiv.org/abs/1911.03162)


## geometric interpretation of Yang-Mills matrix models:

$$S = \text{Tr}([Y^a, Y^b][Y_a, Y_b] + \dots)$$

solution  $Y^a$  (**matrix configuration**) interpreted as

$$Y^a \sim y^a: \mathcal{M} \rightarrow \mathbb{R}^D$$

$\mathcal{M}$  ... **symplectic manifold** (“brane”)



$Y^a$  generates algebra of functions  $\text{End}(\mathcal{H}) \sim \mathcal{C}(\mathcal{M})$

semi-classical regime:

$$[\Phi, \Psi] \sim i\{\phi, \psi\}$$

$$S \sim \int_{\mathcal{M}} \rho_M(-\{y^a, y^b\}\{y_a, y_b\} + \dots), \quad \rho_M = \sqrt{|\theta_{\mu\nu}^{-1}|}$$

~ higher-dim. Poisson-sigma model !

noncommutativity of  $\mathcal{M}$  is essential:

- commutative solutions exist, measure 0  
(correspond to Lagrangian submanifolds, hidden double dimensions)
- generic matrix configurations = quantized symplectic spaces

HS 2009.03400, Ishiki 1503.01230

# The effective metric in M.M.

general rule: write kinetic term in semi-class. form, extract eff. metric

eff. action for fluctuations  $Y^a + \mathcal{A}^a$  around background (solution)  $Y^a$ :

$$S[Y + \mathcal{A}] = S[Y] - \text{Tr}([Y^b, \mathcal{A}^a][Y^b, \mathcal{A}^a] + (\text{gauge} - \text{fix etc.}))$$

simplest: transversal fluctuations = scalar fields  $\phi \in \text{End}(\mathcal{H})$

$$\begin{aligned} S[\phi] &= -\text{Tr} \eta_{ab} [Y^a, \phi] [Y^b, \phi] \\ &\sim \int \rho_M \eta_{ab} E^{a\mu} \partial_\mu \phi E^{b\nu} \partial_\nu \phi \sim \int \sqrt{|G|} G^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \end{aligned}$$

semi-classical:

$$-i[Y^a, \phi] \sim \{Y^a, \phi\} = E^{a\mu} \partial_\mu \phi$$

eff. vielbein (frame)

$$E^a := \{Y^a, \cdot\}, \quad E^{a\mu} := \{Y^a, x^\mu\}$$

governs **all** fluctuations in M.M.

## eff. metric

$$G^{\mu\nu} = \rho^{-2} \eta_{ab} E^{a\mu} E^{b\nu} = \rho^{-2} \gamma^{\mu\nu}$$

$$\rho^2 = \rho_M \sqrt{|\gamma^{\mu\nu}|} \quad \dots \text{dilaton}$$

governs **all** fluctuations in M.M.  
universal metric  $\Rightarrow$  **gravity** !?

HS 1003.4134 ff

can show:  $\square = -\{Y^a, \{Y_a, \cdot\}\} = \rho^2 \square_G \quad \dots$  Matrix Laplacian

issues:

- frame  $E^a = \{Y^a, \cdot\}$  ... not enough dof on 4D  $\mathcal{M}$  (... ?)
- $\theta^{\mu\nu}$  explicitly breaks Lorentz invariance
- class. M.M.: no Einstein equations ("pre-gravity")  
but: quantum effects  $\rightarrow$  **induced gravity** (below, cf. Sakharov)

## eff. metric

$$G^{\mu\nu} = \rho^{-2} \eta_{ab} E^{a\mu} E^{b\nu} = \rho^{-2} \gamma^{\mu\nu}$$

$$\rho^2 = \rho_M \sqrt{|\gamma^{\mu\nu}|} \quad \dots \text{dilaton}$$

governs **all** fluctuations in M.M.  
universal metric  $\Rightarrow$  **gravity** !?

HS 1003.4134 ff

can show:  $\square = -\{Y^a, \{Y_a, \cdot\}\} = \rho^2 \square_G \quad \dots$  Matrix Laplacian

issues:

- frame  $E^a = \{Y^a, \cdot\}$  ... not enough dof on 4D  $\mathcal{M}$  (... ?)
- $\theta^{\mu\nu}$  explicitly breaks Lorentz invariance
- class. M.M.: no Einstein equations ("pre-gravity")  
but: quantum effects  $\rightarrow$  **induced gravity** (below, cf. **Sakharov**)



# 4D covariant quantum spaces & $\mathfrak{h}_5$

key step:

symplectic  $S^2$ - bundle over space(time)  $\mathcal{M}$  (equivariant)

- $\langle \theta^{\mu\nu} = 0 \rangle_{\mathcal{M}}$  !
- price to pay: higher-spin theory
- **vol.-preserving diffeos** on  $\mathcal{M} \subset$  higher-dim symplectomorphisms

main examples:

- prototype: fuzzy  $S_N^4$   
 Grosse-Klimcik-Presnajder; Castellino-Lee-Taylor; Medina-o'Connor;  
 Ramgoolam; Kimura; Abe; Karabail-Nair; Zhang-Hu; HS
- noncompact  $H_n^4$       Hasebe 1207.1968 , M. Sperling, HS 1806.05907
- projection  $\rightarrow$  cosmological space-time  $\mathcal{M}_n^{3,1}$   
 HS, 1710.11495, M. Sperling, HS 1901.03522, ff.

# Euclidean fuzzy hyperboloid $H_n^4$ ( $=EAdS_n^4$ )

$\mathcal{M}^{ab}$  ... hermitian generators of  $\mathfrak{so}(4, 2)$

$$[\mathcal{M}_{ab}, \mathcal{M}_{cd}] = i(\eta_{ac}\mathcal{M}_{bd} - \eta_{ad}\mathcal{M}_{bc} - \eta_{bc}\mathcal{M}_{ad} + \eta_{bd}\mathcal{M}_{ac}) .$$

choose “short” discrete unitary irreps  $\mathcal{H}_n$  (“minireps”, doubletons)


- irreps under  $\mathfrak{so}(4, 1)$
- positive discrete spectrum  $\text{spec}(\mathcal{M}^{05}) = \{E_0, E_0 + 1, \dots\}$

5 generators

$$X^a := r\mathcal{M}^{a5}, \quad a = 0, \dots, 4$$

satisfy

$$\begin{aligned} \eta_{ab}X^aX^b &= X^iX^i - X^0X^0 = -R^2\mathbf{1} \\ R^2 &= r^2(n^2 - 4) \end{aligned}$$



hyperboloid  $H^4 \subset \mathbb{R}^{1,4}$ , covariant under  $SO(4, 1)$

# Euclidean fuzzy hyperboloid $H_n^4$ (= $EAdS_n^4$ )

$\mathcal{M}^{ab}$  ... hermitian generators of  $\mathfrak{so}(4, 2)$

$$[\mathcal{M}_{ab}, \mathcal{M}_{cd}] = i(\eta_{ac}\mathcal{M}_{bd} - \eta_{ad}\mathcal{M}_{bc} - \eta_{bc}\mathcal{M}_{ad} + \eta_{bd}\mathcal{M}_{ac}) .$$

choose “short” discrete unitary irreps  $\mathcal{H}_n$  (“minireps”, doubletons)

- irreps under  $\mathfrak{so}(4, 1)$
- positive discrete spectrum  $\text{spec}(\mathcal{M}^{05}) = \{E_0, E_0 + 1, \dots\}$


5 generators

$$X^a := r\mathcal{M}^{a5}, \quad a = 0, \dots, 4$$

satisfy

$$\begin{aligned} \eta_{ab}X^aX^b &= X^iX^i - X^0X^0 = -R^2\mathbf{1} \\ R^2 &= r^2(n^2 - 4) \end{aligned}$$

hyperboloid  $H^4 \subset \mathbb{R}^{1,4}$ , covariant under  $SO(4, 1)$



claim:


$$H_n^4 = \text{quantized } \mathbb{C}P^{1,2} = S^2 \text{-bundle over } H^4$$

best seen from oscillator construction:

4 bosonic oscillators  $[\psi_\alpha, \bar{\psi}^\beta] = \delta_\alpha^\beta$

$$\mathcal{M}^{ab} = \bar{\psi} \Sigma^{ab} \psi, \quad \gamma_0 = \text{diag}(1, 1, -1, -1)$$

$$X^a = \bar{\psi} \gamma^a \psi$$



$End(\mathcal{H}_n) \cong$  functions on  $\mathbb{C}P^{1,2} \cong$  functions on  $H^4 \otimes$  harmonics on  $S_n^2$

would-be KK modes  $\rightarrow$  higher spin modes

$$End(\mathcal{H}_n) \cong \bigoplus_{s=0}^n \mathcal{C}^s$$

matrix model  $\rightarrow$  higher spin gauge theory, truncated at  $n$

# $\mathcal{M}^{3,1}$ FLRW quantum space-time

generated by

$$\bar{Y}^\mu := \frac{1}{R} \mathcal{M}^{\mu 4}, \quad \square \bar{Y}^\mu = \frac{3}{R^2} \bar{Y}^\mu, \quad \mu = 0, \dots, 3$$

(= solution of IKKT model with mass term)


eff. metric  $\square = [\bar{Y}^\mu, [\bar{Y}_\mu, \cdot]] \cong \rho^2 \square_G$

→ Lorentzian effective metric:

$$ds_G^2 = -dt^2 + a(t)^2 d\Sigma^2 \dots \text{FLRW space-time } \mathcal{M}^{3,1}$$

= projection of  $H_n^4 \subset \mathbb{R}^{1,4} \xrightarrow{\pi} \mathbb{R}^{1,3}$ .

manifest  $SO(3, 1)$ , Big Bounce



# fluctuations & $\mathfrak{h}_5$ gauge fields on $\mathcal{M}^{3,1}$

add **fluctuations** to  $\mathcal{M}^{3,1}$  background

$$Y^\mu = \bar{Y}^\mu + \mathcal{A}^\mu$$

expand action to second order in  $\mathcal{A}_\mu(x, t)$  ...  $\mathfrak{h}_5$ -valued 1-form on  $\mathcal{M}^{3,1}$

$$S[Y] = S[\bar{Y}] + \frac{2}{g^2} \text{Tr} \mathcal{A}_\mu \left( \underbrace{\left( (\square - \frac{3}{R^2}) \delta_\nu^\mu + 2[[\bar{Y}^\mu, \bar{Y}^\nu], \cdot] \right)}_{\mathcal{D}^2} - \underbrace{[[\bar{Y}^\mu, [\bar{Y}^\nu, \cdot]]]}_{g.f.} \right) \mathcal{A}_\nu$$

eigenmodes of  $\mathcal{D}^2$ :

M. Sperling, HS: 1901.03522

- 4 towers of **off-shell modes** for each  $s > 0$
- 4 towers of **on-shell modes** for each  $s > 0$ , massless universal propagation  $\square \sim \square_G$
- 2 spin 0 modes (+ tower of exceptional spin 0 modes)

gauge-fixing  $\{t^\mu, \mathcal{A}_\mu\} = 0$

physical Hilbert space

$$\mathcal{H}_{\text{phys}} = \{\mathcal{D}^2 \mathcal{A} = 0, \mathcal{A} \text{ gauge fixed}\} / \{\text{pure gauge}\}$$

results:

- generically **2 physical modes**  $\square \phi^{(s)} = 0$  for each  $s \geq 1$   
would-be massive,  $m^2 = 0$  (+ exceptional spin 0 modes)
- spin 2 Ricci-flat metric fluctuations
- **no ghosts** ( $t^\mu$  is space-like!) HS 1910.00839  
no tachyons
- same propagation for all modes

... consistent higher-spin gauge theory

# nonlinear regime: frame

any background  $Y^{\dot{\alpha}}$  defines  $\mathfrak{h}_S$  - valued **frame**

fundamental object: **frame**

$$E^{\dot{\alpha}\mu} = \{Y^{\dot{\alpha}}, X^\mu\}$$

$$\text{divergence constraint } \nabla_\nu(\rho^{-2} E_{\dot{\alpha}}{}^\nu) = 0$$

(Jacobi)

**no local Lorentz transformation of the frame!** (diffeo  $\checkmark$ )

eff. metric:

$$G^{\mu\nu} = \rho^{-2} \eta^{\dot{\alpha}\dot{\beta}} E_{\dot{\alpha}}{}^\mu E_{\dot{\beta}}{}^\nu \quad (\mathfrak{h}_S - \text{valued})$$

conjecture:

for any metric, there is a frame  $E^{\dot{\alpha}}$  such that the divergence-constraint is satisfied, thereby fixing  $\rho, \tilde{\rho}$



# gauge transformations and ( $\mathfrak{h}_5$ -valued) diffeos

scalar fields:

$$\delta_{\Lambda}\phi = \{\Lambda, \phi\} = \xi^{\mu}\partial_{\mu}\phi = \mathcal{L}_{\xi}\phi, \quad \xi^{\mu} = \{\Lambda, x^{\mu}\}$$

... push-forward of Hamiltonian VF (symplectomorphisms) to  $\mathcal{M}$  by bundle projection

frame:

$$\delta_{\Lambda}Y_{\dot{\alpha}} = \{\Lambda, Y_{\dot{\alpha}}\}$$

$$(\delta_{\Lambda}E_{\dot{\alpha}})\phi = \{\Lambda, \{Y_{\dot{\alpha}}, \phi\}\} - \{Y_{\dot{\alpha}}, \{\Lambda, \phi\}\} = (\mathcal{L}_{\xi}E_{\dot{\alpha}})\phi \quad (\text{Jacobi})$$

hence

$$\delta_{\Lambda}E_{\dot{\alpha}}^{\mu} = \mathcal{L}_{\xi}E_{\dot{\alpha}}^{\mu}, \quad \delta_{\Lambda}G^{\mu\nu} = \mathcal{L}_{\xi}G^{\mu\nu}$$

diffeos from NC gauge trafos

( $\mathfrak{h}_5$ -valued) **Weitzenböck connection**:

$$\nabla^{(W)} E_{\dot{\alpha}} = 0 \quad (\text{Weitzenböck}) \quad \Rightarrow \quad \nabla^{(W)} G^{\mu\nu} = 0$$

flat but ( $\mathfrak{h}_5$  - valued) **torsion**:

$$T_{\dot{\alpha}\dot{\beta}} \equiv T[E_{\dot{\alpha}}, E_{\dot{\beta}}] = \nabla_{\dot{\alpha}} E_{\dot{\beta}} - \nabla_{\dot{\beta}} E_{\dot{\alpha}} - [E_{\dot{\alpha}}, E_{\dot{\beta}}]$$

can show:

$$T_{\dot{\alpha}\dot{\beta}}{}^{\mu} = \{\hat{\Theta}_{\dot{\alpha}\dot{\beta}}, x^{\mu}\}, \quad \hat{\Theta}_{\dot{\alpha}\dot{\beta}} := -\{Y_{\dot{\alpha}}, Y_{\dot{\beta}}\}$$

$$T_{\dot{\alpha}} = dE_{\dot{\alpha}}, \quad E_{\dot{\alpha}} = E_{\mu\dot{\alpha}} dx^{\mu} \quad \dots \text{coframe}$$

torsion tensor encodes field strength of the NC gauge theory

HS arXiv:2002.02742 , Langmann Szabo hep-th/0105094

# classical dynamics of geometry: pre-gravity

matrix model eom:

$$\{Y^{\dot{\alpha}}, \hat{\Theta}_{\dot{\alpha}\dot{\beta}}\} = m^2 Y_{\dot{\beta}}$$

can recast as  $\nabla_{\nu}^{(W)} T^{\nu}_{\rho\mu} + T_{\nu}^{\sigma}_{\mu} T_{\sigma\rho}^{\nu} = m^2 \rho^{-2} G_{\rho\mu}$

HS arXiv:2002.02742 cf. Furuta, Hanada, Kawai, Kimura hep-th/0611093

→ eom for frame:

$$d(\rho^2 \star T^{\dot{\alpha}}) = d\tilde{\rho} \wedge T^{\dot{\alpha}} + m^2 \star E^{\dot{\alpha}} .$$

tot. AS part:  $\epsilon^{\mu\nu\rho\sigma} T_{\mu\nu\rho} \stackrel{eom}{\sim} \partial^{\sigma} \tilde{\rho} \dots$  axion

contraction:  $T_{\mu\nu}^{\mu} \sim \rho^{-1} \partial_{\nu} \rho \dots$  dilaton

fully covariant form of matrix model eom for frame, axion & dilaton

S. Fredenhagen, HS arXiv: 2101.07297

## pre-gravity from classical matrix model:

( $\mathfrak{h}_5$ -extended) theory of dynamical geometry, similar to gravity  
differs from GR at non-lin level

- univ. metric, gravitons, lin. Schwarzschild etc. recovered
- extra dof: dilaton, axion,  $\mathfrak{h}_5$ ; massive graviton modes (?)
- bare action:  $S \sim \int \frac{1}{g^2} \Theta_{\dot{\alpha}\dot{\beta}} \Theta^{\dot{\alpha}\dot{\beta}}$  ... 2 derivatives **less** than

$$\int d^4x \sqrt{|G|} \mathcal{R} = \int d^4x \sqrt{|G|} \left( -\frac{3}{4} T_\nu T_\mu G^{\mu\nu} - \frac{7}{8} T^\mu{}_{\sigma\rho} T_{\mu\sigma'}{}^\rho G^{\sigma\sigma'} \right)$$

where  $T^{\dot{\alpha}\dot{\beta}\mu} = \{\Theta^{\dot{\alpha}\dot{\beta}}, x^\mu\} \sim \partial \Theta^{\dot{\alpha}\dot{\beta}}$

S. Fredenhagen, H.S. arxiv:2101.07297

⇒ **different** from GR, expected to dominate on **large scales**  
**good for quantization!**

- reasonable cosmology without any fine-tuning

# 1-loop effective action and gravity

SUSY model  $\rightarrow$  quantum effects mild:

gravity = quantum effect on quantum space-time in MM

- Einstein-Hilbert action (+ extra) arises in the 1-loop effective action on  $\mathcal{M}^{3,1}$  space-time (cf. Sakharov '67)

$$\Gamma_{1\text{-loop}} \ni \int_{\mathcal{M}} T_{\nu\lambda}{}^{\mu} T_{\nu\lambda}{}^{\mu} + \dots \sim \int_{\mathcal{M}} d^4x \sqrt{G} m_{\mathcal{K}}^2 \mathcal{R}[G] + \dots$$

requires presence of fuzzy extra dimensions  $\mathcal{K}$

- different from bulk IIB sugra = short-range  $r^{-8}$  correction! (also 1-loop)
- gravity action on brane = IIB sugra **interaction** of  $\mathcal{M}$  with  $\mathcal{K}$

nonperturbative quantization:

$$Z = \int dY d\Psi e^{iS[Y, \Psi]}, \quad S = S_{\text{IKKT}} + i\varepsilon Y^a Y^b \delta_{ab}$$

cf. next workshop (Nishimura, Tsuchiya,...)

1-loop effective action

$$e^{i\Gamma_{1\text{-loop}}[Y]} = \int_{1 \text{ loop}} d\mathcal{A} d\Psi e^{iS[Y+\mathcal{A}, \Psi]}$$

$$\begin{aligned} \Gamma_{1\text{loop}}[Y] &= \frac{1}{2} \text{Tr} \left( \log(\square - M_{ab}[\Theta^{ab}, \cdot]) - \frac{1}{2} \log(\square - M_{ab}^{(\psi)}[\Theta^{ab}, \cdot]) - 2 \log(\square) \right) \\ &= \frac{1}{2} \text{Tr} \left( \sum_{n=4}^{\infty} \frac{1}{n} \left( (\square^{-1} M_{ab}[\Theta^{ab}, \cdot])^n - \frac{1}{2} (\square^{-1} M_{ab}^{(\psi)}[\Theta^{ab}, \cdot])^n \right) \right) \end{aligned}$$

(max. **SUSY** !) where

$$i\Theta^{ab} = -[Y^a, Y^b]$$

$M_{ab}^{(\psi)}$  ...spinorial generators of  $so(5)$

$M_{ab}$  ...vector generators


UV-finite on 4D backgrounds due to max. SUSY

evaluate **trace** use **string state formalism**

$$\text{Tr}_{\text{End}(\mathcal{H})} \mathcal{O} = \frac{1}{(2\pi)^m} \int_{\mathcal{M} \times \mathcal{M}} dx dy \langle x | \mathcal{O} | y \rangle$$

string states:

$$\boxed{\begin{matrix} |x \\ |y \end{matrix} \rangle := |x\rangle\langle y|} \in \text{End}(\mathcal{H})$$



$|x\rangle$  ... coherent state on  $\mathcal{M}$

$\approx$  diagonalize kinetic operators:

$$\begin{aligned} \square^{-1} |x \\ |y \rangle &\approx \frac{1}{|x-y|^2 + 2\Delta^2} |x \\ |y \rangle \\ \square^{-1} [\Theta^{ab}, \cdot] |x \\ |y \rangle &\approx \frac{1}{|x-y|^2 + 2\Delta^2} \underbrace{(\Theta^{ab}(y) - \Theta^{ab}(x))}_{\delta\Theta^{ab}} |x \\ |y \rangle \end{aligned}$$

can evaluate

$$\begin{aligned}\Gamma_{\text{loop};4}[Y] &= \frac{1}{8} \text{Tr} \left( (\square^{-1}(M_{ab}[\Theta^{ab}, \cdot])^4 - \frac{1}{2}(\square^{-1}M_{ab}^{(\psi)}[\Theta^{ab}, \cdot])^4) \right) \\ &= \frac{1}{4} \frac{1}{(2\pi)^m} \int_{\mathcal{M} \times \mathcal{M}} dx dy \frac{3S_4[\delta\Theta(x,y)]}{(|x-y|^2 + 2\Delta^2)^4}\end{aligned}$$

where

$$-S_4[\delta\Theta] = 4 \text{tr}(\delta\Theta\delta\Theta\delta\Theta\delta\Theta) - (\text{tr}\delta\Theta\delta\Theta)^2$$

note:

- UV regime = long string regime (  $\rightarrow$  UV/IR mixing, non-local)  
(cf. talk by [Tekel](#) tomorrow)
- UV-finite only in maximally SUSY model  $\rightarrow$   
IIB supergravity in  $\mathbb{R}^{9,1}$ ,  $\sim r^{-8}$  interaction
- short-distance regime requires refined analysis:



string states as localized Gaussian wave-packets:

local isometric mapping  $\text{End}(\mathcal{H})_{loc} \rightarrow \mathcal{C}_{IR}(\mathcal{M})$  :

$$e^{\frac{i}{2} k^a \theta_{ab}^{-1} y^b} \left| y \pm \frac{k}{2} \right\rangle =: \Psi_{\tilde{k};y} \cong \psi_{\tilde{k};y}(x) = \frac{2}{\pi L_{NC}^2} e^{i\tilde{k}x} e^{-|x-y|_g^2} .$$


(for almost-Kähler  $(\mathcal{M}, \theta^{-1}, g)$ ), size  $\sim L_{NC}$

$$\tilde{k}_\mu = \theta_{\mu\nu}^{-1} k^\nu$$

H.S., J. Tekel arxiv:21xx.xxxxx

semi-classical wavepackets:

$$\Psi_{k;y}^{(L)} := \int d^4z e^{-|y-z|^2/L^2} \Psi_{k;z} \cong e^{ikx} e^{-|x-y|^2/L^2} =: \psi_{k;y}^{(L)}$$



locally diagonalizes kinetic operators in IR:

$$\square \Psi_{k;y}^{(L)} \approx \gamma^{\mu\nu}(x) k_\mu k_\nu \Psi_{k;y}^{(L)}$$

$$[\theta^{ab}, \Psi_{k;y}^{(L)}] \approx i \{ \theta^{ab}, \psi_{k;y}^{(L)} \} \approx - \{ \theta^{ab}, x^\mu \} k_\mu \psi_{k;y}^{(L)}$$


## Trace formula for UV-finite traces on NC spaces:

$$\text{Tr} \mathcal{O} = \frac{1}{(2\pi)^m} \int_{\mathcal{M} \times \mathcal{M}} \Omega_x \Omega_y \langle x | \mathcal{O} | x \rangle \approx \frac{1}{(2\pi)^m} \int_{\mathcal{M}} \sqrt{G} dx \int \frac{1}{\sqrt{G}} dk \langle \Psi_{k,x}^{(L)}, \mathcal{O} \Psi_{k,x}^{(L)} \rangle$$

apply this to  $\mathcal{O} = (\square^{-1}[\Theta^{ab}, \cdot])^4$  in 1-loop eff. action

a priori: 4-derivative action ☺

however: assume **fuzzy extra dimensions**  $\mathcal{M} \times \mathcal{K}$   
from 6 transversal directions  $\langle \phi^i \rangle = \lambda^i \neq 0$



( $\rightarrow$  interesting gauge theory, cf. H.S., Zahn arxiv:1409.1440, Chatzistavrakidis Zoupanos 1107.0265 ...)

mixed term  $(\delta\Theta^{\alpha\beta} \delta\Theta^{\alpha\beta}) (\delta\Theta^{ij} \delta\Theta^{ij})$  leads to induced E-H action ☺

induced E-H action:

$$\begin{aligned} \{\theta^{\alpha\beta}, \{\theta^{\alpha\beta}, \psi_{k;y}\}\} &\approx -\{\theta^{\alpha\beta}, x^\mu\} \{\theta^{\alpha\beta}, x^\nu\} k_\mu k_\nu \psi_{k;y} \\ &= -T^{\alpha\beta\mu} k_\mu T^{\alpha\beta\nu} k_\nu \psi_{k;y}, \quad k > \frac{1}{L}. \end{aligned}$$

recall torsion  $T^{\alpha\beta\mu} = \{\theta^{\alpha\beta}, x^\mu\}$

$$\begin{aligned} \Gamma_{\text{1loop}} = i\text{Tr} \left( \frac{V_{4,\text{mix}}}{(\square - i\varepsilon)^4} \right) &\sim \int_{\mathcal{M}} d^4x \sum_{lm} C_{lm}^2 \int d^4k \frac{T^{\alpha\beta\mu} k_\mu T^{\alpha\beta\nu} k_\nu}{(k \cdot k + m_{lm}^2 - i\varepsilon)^4} \\ &\sim - \int_{\mathcal{M}} d^4x \sqrt{G} m_{\mathcal{K}}^2 T^\rho_{\sigma\mu} T_{\rho'}^{\sigma\mu} G^{\mu\mu'} \\ &\sim \int d^4x \sqrt{G} m_{\mathcal{K}}^2 \left( 8\mathcal{R}[G] + 6T_\nu T_\mu G^{\mu\nu} \right) \end{aligned}$$

$m_{\mathcal{K}}^2$  ... Kaluza-Klein mass scale on  $\mathcal{K}$

bottom line:

(to be published)

- $\Gamma_{\text{loop}}$  includes Einstein-Hilbert action

$$\frac{1}{16\pi G_N} \sim c_{\mathcal{K}}^2 m_{\mathcal{K}}^2 \quad \dots \text{ eff. Newton constant}$$

set by Kaluza-Klein mass scale on  $\mathcal{K}$

- vacuum energy

$$\Gamma_{\text{loop}} \ni C \int \Omega \quad \dots \text{ symplectic volume, non-dynamical}$$

**no induced cosm. const !**

- in addition to matrix model action  $S \sim \int [Y, Y][Y, Y]$ , should dominate extreme IR (cosm.)
- attractive potential between  $\mathcal{K}$  and cosm.  $\mathcal{M}$
- + lots of other stuff (axion, dilaton, ...  $\hbar$ s), to be understood

# summary

gravity arises as quantum effect on 3+1-dim. quantum space-time in the maximally SUSY IKKT matrix model

- "pre-gravity", suitable for quantization
- covariant quantum spaces = twisted  $S^2$  bundles over  $\mathcal{M}^{3,1}$   
→ **higher spin gauge theory**  
Lorentz invariance partially manifest
- quantization → **induced Einstein-Hilbert action**, no c.c. problem
- expect cross-over GR ↔ cosm. background (class.)
- new physics (**axion, dilaton,  $\mathfrak{h}_5$  ...**)  
lots of things to be clarified !

# linearized Schwarzschild solution

HS 1905.07255

Ricci-flat "scalar" metric perturbation

from  $\mathcal{A}^{(-)}[D^+ D^+ \phi]$

$$ds^2 = (G_{\mu\nu} - h_{\mu\nu}) dy^\mu dy^\nu = -dt^2 + a(t)^2 d\Sigma^2 + \phi' (dt^2 + a(t)^2 d\Sigma^2)$$

$$\phi' \sim \frac{M(t)}{2r} \frac{1}{a(t)}$$

$\approx$  lin. Schwarzschild (Vittie) solution on FRW, eff. mass  $M(t) \sim \frac{M_0}{a(t)}$

more generally for any quasi-static lin. solution in GR

# non-linear solution of class. equations

solve  $d(\rho^2 \star T^{\dot{\alpha}}) = d\tilde{\rho} \wedge T^{\dot{\alpha}} + m^2 \star E^{\dot{\alpha}} \Leftrightarrow$

$$\nabla_{(G)}^{\nu}(\rho^2 T_{\nu\rho}^{\dot{\alpha}}) = \frac{1}{2} \sqrt{|G|}^{-1} \varepsilon^{\nu\rho'\sigma\mu} G_{\rho\rho'} \partial_{\mu} \tilde{\rho} T_{\nu\sigma'}^{\dot{\alpha}} + m^2 E_{\rho}^{\dot{\alpha}} .$$

spherically symmetric static solutions for frame:

simplest solution:

S. Fredenhagen, HS arXiv: 2101.07297

$$ds_G^2 = -\frac{c_1 b_0^{-2}}{(1+\frac{M}{r})} dt^2 + c_1 \left(1 + \frac{M}{r}\right) (dr^2 + r^2 d\Omega^2)$$

$$\rho^2 = c_1 b_0^{-2} \left(1 + \frac{M}{r}\right) .$$

linearized Schwarzschild, deviates at nonlin. level

most general spherical solution:

Y. Asano, HS arXiv: 21xx.xxxxx