

Roles of dark Higgs Boson in DM physics and Higgs inflation


Pyungwon Ko (KIAS)

CORFU 2022
Aug 28-Sep 8 (2022)

Contents

- Motivations
- Higgs portal DM : EFT vs. UV Completions
 - DM searches @ high energy colliders, $\Gamma_{\text{inv}}(H \rightarrow VV)$ for $m_V \rightarrow 0$,
 - GC γ -ray Excess, etc. [(dark) gauge invariance and unitarity]
- Inelastic DM for light WIMP (cf. XENON1T Excess)
- $U(1)_{L_\mu - L_\tau}$ -charged DM : Z' only vs. $Z' + \phi$
- Higgs-Portal Assisted Higgs Inflation
- Summary

Motivations

- Existence of DM established from astro/cosmo observations
- DM-DM, DM-SM force carried by some mediators
- Dark photon : the most common. Usually the origin of its mass is ignored (assuming Stueckelberg mechanism)
- Dark Higgs : a kind of the must for massive dark photon within perturbation  Main theme of this talk
- DM phenomenology can change drastically !

Higgs Portal DM : EFT vs. UV completions

$$\Gamma_{\text{inv}}(H \rightarrow VV) \text{ for } m_V \rightarrow 0$$

arXiv: 2112.11983, PRD 105 (2022) 015007, with S. Baek, W.I. Park
And references therein by P. Ko et al

Higgs portal DM models

All invariant
under ad hoc
Z2 symmetry

$$\mathcal{L}_{\text{scalar}} = \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_S^2 S^2 - \frac{\lambda_{HS}}{2} H^\dagger H S^2 - \frac{\lambda_S}{4} S^4$$

$$\mathcal{L}_{\text{fermion}} = \bar{\psi} [i\gamma \cdot \partial - m_\psi] \psi - \frac{\lambda_{H\psi}}{\Lambda} H^\dagger H \bar{\psi} \psi$$

$$\mathcal{L}_{\text{vector}} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + \frac{1}{2} m_V^2 V_\mu V^\mu + \frac{1}{4} \lambda_V (V_\mu V^\mu)^2 + \frac{1}{2} \lambda_{HV} H^\dagger H V_\mu V^\mu.$$

arXiv:1112.3299, ... 1402.6287, etc.




FIG. 1. Scalar Higgs-portal parameter space allowed by WMAP (between the solid red curves), XENON100 and $\text{Br}^{\text{inv}} = 10\%$ for $m_h = 125$ GeV. Shown also are the prospects for XENON upgrades.




FIG. 2. Same as Fig. 1 for vector DM particles.




FIG. 3. Same as in Fig.1 for fermion DM; λ_{hff}/Λ is in GeV^{-1} .

Higgs portal DM as examples

$$\mathcal{L}_{\text{scalar}} = \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_S^2 S^2 - \frac{\lambda_{HS}}{2} H^\dagger H S^2 - \frac{\lambda_S}{4} S^4$$

$$\mathcal{L}_{\text{fermion}} = \bar{\psi} [i\gamma \cdot \partial - m_\psi] \psi - \frac{\lambda_{H\psi}}{\Lambda} H^\dagger H \bar{\psi} \psi$$

$$\mathcal{L}_{\text{vector}} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + \frac{1}{2} m_V^2 V_\mu V^\mu + \frac{1}{4} \lambda_V (V_\mu V^\mu)^2 + \frac{1}{2} \lambda_{HV} H^\dagger H V_\mu V^\mu.$$

All invariant
under ad hoc
Z2 symmetry

arXiv:1112.3299, ... 1402.6287, etc. And Revived recent papers

**We need to include dark Higgs or singlet scalar
to get renormalizable/unitary models
for Higgs portal singlet fermion or vector DM
[NB: UV Completions : Not unique]**

Models for HP SFDM & VDM

UV Completion of HP Singlet Fermion DM (SFDM)

$$\begin{aligned}\mathcal{L} = & \mathcal{L}_{\text{SM}} - \mu_{HS} S H^\dagger H - \frac{\lambda_{HS}}{2} S^2 H^\dagger H \\ & + \frac{1}{2} (\partial_\mu S \partial^\mu S - m_S^2 S^2) - \mu'_S S - \frac{\mu''_S}{3} S^3 - \frac{\lambda_S}{4} S^4 \\ & + \bar{\psi} (i \not{\partial} - m_{\psi_0}) \psi - \lambda S \bar{\psi} \psi\end{aligned}$$

UV Completion of HP VDM

$$\begin{aligned}\mathcal{L}_{VDM} = & -\frac{1}{4} X_{\mu\nu} X^{\mu\nu} + (D_\mu \Phi)^\dagger (D^\mu \Phi) - \frac{\lambda_\Phi}{4} \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2} \right)^2 \\ & - \lambda_{H\Phi} \left(H^\dagger H - \frac{v_H^2}{2} \right) \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2} \right),\end{aligned}$$

- The simplest UV completions in terms of # of new d.o.f.
- At least, 2 more parameters, $(m_\phi, \sin \alpha)$ for DM physics


UV Completion for HP FDM

Baek, Ko, Park, arXiv:1112.1847

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_{\text{SM}} - \mu_{HS} S H^\dagger H - \frac{\lambda_{HS}}{2} S^2 H^\dagger H \\ & + \frac{1}{2} (\partial_\mu S \partial^\mu S - m_S^2 S^2) - \mu_S^3 S - \frac{\mu'_S}{3} S^3 - \frac{\lambda_S}{4} S^4 \\ & + \bar{\psi} (i \not{\partial} - m_{\psi_0}) \psi - \lambda S \bar{\psi} \psi \end{aligned}$$

mixing

invisible
decay



Production and decay rates are suppressed relative to SM.

Higgs-Singlet Mixing

- Mixing and Eigenstates of Higgs-like bosons

$$\mu_H^2 = \lambda_H v_H^2 + \mu_{HS} v_S + \frac{1}{2} \lambda_{HS} v_S^2,$$

$$m_S^2 = -\frac{\mu_S^3}{v_S} - \mu'_S v_S - \lambda_S v_S^2 - \frac{\mu_{HS} v_H^2}{2v_S} - \frac{1}{2} \lambda_{HS} v_H^2,$$

at vacuum

$$M_{\text{Higgs}}^2 \equiv \begin{pmatrix} m_{hh}^2 & m_{hs}^2 \\ m_{hs}^2 & m_{ss}^2 \end{pmatrix} \equiv \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$$

$$H_1 = h \cos \alpha - s \sin \alpha,$$

$$H_2 = h \sin \alpha + s \cos \alpha.$$





Mixing of Higgs and singlet

Constraints

- Dark matter to nucleon cross section (constraint)

$$\sigma_p \approx \frac{1}{\pi} \mu^2 \lambda_p^2 \simeq 2.7 \times 10^{-2} \frac{m_p^2}{\pi} \left| \left(\frac{m_p}{v} \right) \lambda \sin \alpha \cos \alpha \left(\frac{1}{m_1^2} - \frac{1}{m_2^2} \right) \right|^2$$



Low energy pheno.

- Universal suppression of collider SM signals


[See I I 12.1847, Seungwon Baek, P. Ko & WIP]

- If “ $m_h > 2 m_\phi$ ”, non-SM Higgs decay!



- Tree-level shift of $\lambda_{H,SM}$ (& loop correction)

$$\lambda_{\Phi H} \Rightarrow \lambda_H = \left[1 + \left(\frac{m_\phi^2}{m_h^2} - 1 \right) \sin^2 \alpha \right] \lambda_H^{SM}$$

➔ If “ $m_\phi > m_h$ ”, vacuum instability can be cured.



[G. Degrassi et al., 1205.6497]



[S. Baek, P. Ko, WIP & E. Senaha, JHEP(2012)]

UV Completion of HP VDM

[S Baek, P Ko, WI Park, E Senaha, arXiv:1212.2131 (JHEP)]

$$\mathcal{L}_{VDM} = -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} + (D_\mu\Phi)^\dagger(D^\mu\Phi) - \frac{\lambda_\Phi}{4}\left(\Phi^\dagger\Phi - \frac{v_\Phi^2}{2}\right)^2 - \lambda_{H\Phi}\left(H^\dagger H - \frac{v_H^2}{2}\right)\left(\Phi^\dagger\Phi - \frac{v_\Phi^2}{2}\right),$$

$X_\mu \equiv V_\mu$ here

$$\Phi(x) = (v_\phi + \phi(x))/\sqrt{2}$$

- There appear a new singlet scalar (**dark Higgs**) $\phi(x)$ from $\Phi(x)$, which mixes with the SM Higgs boson through Higgs portal interaction ($\lambda_{H\Phi}$ term)
- The effects must be similar to the singlet scalar in the fermion CDM model, and generically true in the DM with dark gauge symmetry
- Can accommodate GeV scale gamma ray excess from GC with $VV \rightarrow \phi\phi$
- **Can modify the Higgs inflation : No tight correlation with top mass**




Figure 6. The scattered plot of σ_p as a function of M_X . The big (small) points (do not) satisfy the WMAP relic density constraint within 3σ , while the red-(black-)colored points gives $r_1 > 0.7$ ($r_1 < 0.7$). The grey region is excluded by the XENON100 experiment. The dashed line denotes the sensitivity of the next XENON experiment, XENON1T.

New scalar (Dark Higgs)
improves EW vacuum stability




Figure 8. The vacuum stability and perturbativity constraints in the α - m_2 plane. We take $m_1 = 125 \text{ GeV}$, $g_X = 0.05$, $M_X = m_2/2$ and $v_\Phi = M_X/(g_X Q_\Phi)$.

Interaction Lagrangians

Scalar DM

$$\mathcal{L}_{\text{SDM}}^{\text{int}} = -h \left(\frac{2m_W^2}{v_h} W_\mu^+ W^{-\mu} + \frac{m_Z^2}{v_h} Z_\mu Z^\mu \right) - \lambda_{HS} v_h h S^2.$$

Singlet FDM

$$\mathcal{L}_{\text{FDM}}^{\text{int}} = - (H_1 \cos \alpha + H_2 \sin \alpha) \left(\sum_f \frac{m_f}{v_h} \bar{f} f - \frac{2m_W^2}{v_h} W_\mu^+ W^{-\mu} - \frac{m_Z^2}{v_h} Z_\mu Z^\mu \right) + g_\chi (H_1 \sin \alpha - H_2 \cos \alpha) \bar{\chi} \chi.$$

Vector DM

$$\mathcal{L}_{\text{VDM}}^{\text{int}} = - (H_1 \cos \alpha + H_2 \sin \alpha) \left(\sum_f \frac{m_f}{v_h} \bar{f} f - \frac{2m_W^2}{v_h} W_\mu^+ W^{-\mu} - \frac{m_Z^2}{v_h} Z_\mu Z^\mu \right) - \frac{1}{2} g_V m_V (H_1 \sin \alpha - H_2 \cos \alpha) V_\mu V^\mu.$$

NB: One can not simply ignore 125 GeV Higgs Boson or singlet scalar by hand, since it would violate gauge invariance and unitarity !

- EFT : Effective operator $\mathcal{L}_{int} = \frac{m_q}{\Lambda_{dd}^3} \bar{q}q\bar{\chi}\chi$
- S.M.: Simple scalar mediator S of

$$\mathcal{L}_{int} = \left(\frac{m_q}{v_H} \sin \alpha \right) S \bar{q}q - \lambda_s \cos \alpha S \bar{\chi}\chi$$
- H.M.: A case where a Higgs is a mediator

$$\mathcal{L}_{int} = - \left(\frac{m_q}{v_H} \cos \alpha \right) H \bar{q}q - \lambda_s \sin \alpha H \bar{\chi}\chi$$
- H.P.: Higgs portal model as in eq. (2).

$$\frac{d\sigma_i}{dm_{\chi\chi}} \propto \left| \frac{\sin 2\alpha g_\chi}{m_{\chi\chi}^2 - m_{H_1}^2 + im_{H_1}\Gamma_{H_1}} - \frac{\sin 2\alpha g_\chi}{m_{\chi\chi}^2 - m_{H_2}^2 + im_{H_2}\Gamma_{H_2}} \right|^2$$

$$\begin{aligned} \text{H.P.} &\longrightarrow \text{H.M.}, \\ &\quad m_{H_2}^2 \gg \hat{s} \\ \text{S.M.} &\longrightarrow \text{EFT}, \\ &\quad m_S^2 \gg \hat{s} \\ \text{H.M.} &\neq \text{EFT}. \end{aligned}$$





FIG. 3: The experimental bounds on M_* at 90% C.L. as a function of m_{H_2} (m_S in S.M. case) in the monojet+ \cancel{E}_T search (upper) and $t\bar{t} + \cancel{E}_T$ search (lower). Each line corresponds to the EFT approach (magenta), S.M. (blue), H.M. (black), and H.P. (red), respectively. The bound of S.M., H.M., and H.P., are expressed in terms of the effective mass M_* through the Eq.(16)-(20). The solid and dashed lines correspond to $m_\chi = 50$ GeV and 400 GeV in each model, respectively.


Collider Implications

$m_h = 125\text{GeV}$, $\text{Br}(H \rightarrow \text{inv}) < 0.51$ at 90% CL



Based on EFTs

$m_h = 125.5\text{GeV}$, $\text{Br}(H \rightarrow \text{inv}) < 0.52$ at 90% CL



- However, in renormalizable unitary models of Higgs portals, **2 more relevant parameters !**

$$\mathcal{L}_{\text{SFDM}} = \bar{\psi}(i\partial - m_\psi - \lambda_\psi S) - \mu_{HS} S H^\dagger H - \frac{\lambda_{HS}}{2} S^2 H^\dagger H$$



$$+ \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_S^2 S^2 - \mu'_S S - \frac{\mu'_S}{3} S^3 - \frac{\lambda_S}{4} S^4.$$

[arXiv: 1405.3530, S. Baek, P. Ko & WIPark, PRD]

$$\sigma_p^{\text{SI}} = (\sigma_p^{\text{SI}})_{\text{EFT}} c_\alpha^4 m_h^4 \mathcal{F}(m_{\text{DM}}, \{m_i\}, v)$$



$$\simeq (\sigma_p^{\text{SI}})_{\text{EFT}} c_\alpha^4 \left(1 - \frac{m_h^2}{m_2^2}\right)^2$$

$$\mathcal{L}_{\text{VDM}} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + D_\mu \Phi^\dagger D^\mu \Phi - \lambda_\Phi \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2}\right)^2 - \lambda_{\Phi H} \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2}\right) \left(H^\dagger H - \frac{v_H^2}{2}\right)$$



Dashed curve
ATLAS, CMS

Interpretation of collider data is **quite model-dependent** in **Higgs portal DMs** and in general



- However, in renormalizable unitary models of Higgs portals, **2 more relevant parameters !**

$$\mathcal{L}_{\text{SFDM}} = \bar{\psi}(i\partial - m_\psi - \lambda_\psi S) - \mu_{HS} S H^\dagger H - \frac{\lambda_{HS}}{2} S^2 H^\dagger H$$





$$+ \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_S^2 S^2 - \mu'_S S - \frac{\mu'_S}{3} S^3 - \frac{\lambda_S}{4} S^4.$$

[arXiv: 1405.3530, S. Baek, P. Ko & WIPark, PRD]

$$\sigma_p^{\text{SI}} = (\sigma_p^{\text{SI}})_{\text{EFT}} c_\alpha^4 m_h^4 \mathcal{F}(m_{\text{DM}}, \{m_i\}, v)$$

$$\simeq (\sigma_p^{\text{SI}})_{\text{EFT}} c_\alpha^4 \left(1 - \frac{m_h^2}{m_2^2}\right)^2$$

$$\mathcal{L}_{\text{VDM}} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + D_\mu \Phi^\dagger D^\mu \Phi - \lambda_\Phi \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2}\right)^2 - \lambda_{\Phi H} \left(\Phi^\dagger \Phi - \frac{v_\Phi^2}{2}\right) \left(H^\dagger H - \frac{v_H^2}{2}\right)$$



Dashed curves: EFT, ATLAS, CMS results


Search for $H \rightarrow$ Dark matter (invisible)

$BR(H \rightarrow \text{invisible}) < 14.5\%$ (obs) (10.3% exp.)

from search with VBF topology


(13% limit when combined with Higgs coupling measurements)

ATLAS Highlight talks
By G.Unal @ICHEP2022, and
By M. Cristinziani @ CORFU2022



Now implemented in the ATLAS results,
But only for VDM, and not for SFDM

arXiv:2202.07953



Invisible H decay into a pair of VDM

[arXiv: 1405.3530, S. Baek, P. Ko & WIPark, PRD]

$$(\Gamma_h^{\text{inv}})_{\text{EFT}} = \frac{\lambda_{VH}^2 v_H^2 m_h^3}{128\pi m_V^4} \times \left(1 - \frac{4m_V^2}{m_h^2} + 12\frac{m_V^4}{m_h^4}\right) \left(1 - \frac{4m_V^2}{m_h^2}\right)^{1/2} \quad (23)$$

Diverge when $m_V \rightarrow 0 !!$

$$m_V \propto g_X Q_\Phi v_\Phi$$

$$\frac{g_X^2}{m_V^2} = \frac{g_X^2}{g_X^2 Q_\Phi^2 v_\Phi^2} \rightarrow \frac{1}{v_\Phi^2} = \text{finite}$$

VS.

$$\Gamma_i^{\text{inv}} = \frac{g_X^2 m_i^3}{32\pi m_V^2} \left(1 - \frac{4m_V^2}{m_i^2} + 12\frac{m_V^4}{m_i^4}\right) \left(1 - \frac{4m_V^2}{m_i^2}\right)^{1/2} \sin^2 \alpha \quad (22)$$

Invisible H decay width : finite for small m_V in unitary/renormalizable model

Two Limits for $m_V \rightarrow 0$

Also see the addendum:
by S Baek, P Ko, WI Park

- $m_V = g_X Q_\Phi v_\Phi$ in the UV completion with dark Higgs boson
- Case I : $g_X \rightarrow 0$ with finite $v_\Phi \neq 0$

$$\frac{g_X^2 Q_\Phi^2}{m_V^2} = \frac{g_X^2 Q_\Phi^2}{g_X^2 Q_\Phi^2 v_\Phi^2} = \frac{1}{v_\Phi^2} = \text{finite.}$$

$$(\Gamma_h^{\text{inv}})_{\text{UV}} = \frac{1}{32\pi} \frac{m_h^3}{v_\Phi^2} \sin^2 \alpha = \Gamma(h \rightarrow a_\Phi a_\Phi)$$

with a_Φ being the NG boson for spontaneously broken global $U(1)_X$

- Case II : $v_\Phi \rightarrow 0$ with finite $g_X \neq 0$

$$\alpha \xrightarrow{v_\Phi \rightarrow 0^+} \frac{2\lambda_{H\Phi} v_\Phi}{\lambda_H v_H}$$


$$\frac{g_X^2 Q_\Phi^2}{m_V^2} \sin^2 \alpha \xrightarrow{v_\Phi \rightarrow 0^+} \frac{4\lambda_{H\Phi}^2}{\lambda_H^2 v_H^2} = \frac{2\lambda_{H\Phi}^2}{\lambda_H m_h^2} = \text{finite,}$$

$$(\Gamma_h^{\text{inv}})_{\text{UV}} \xrightarrow{v_\Phi \rightarrow 0^+} \frac{1}{16\pi} \frac{\lambda_{H\Phi}^2 m_h}{\lambda_H}$$

Therefore $\Gamma(h \rightarrow VV)$ is finite when $m_V \rightarrow 0$ in the UV completions

DM Production @ ILC

P Ko, H Yokoya, arXiv:1603.08802, JHEP





$$t \equiv m_{DD}^2$$

We consider $e^+e^- \rightarrow Z^* \rightarrow ZH_{i=1,2}$
followed by $H_i \rightarrow \bar{\chi}\chi$

$$\frac{d\sigma_{SDM}}{dt} \propto \sigma_{SDM}^{h^*} \times \left| \frac{1}{t - m_h^2 + im_h\Gamma_h} \right|^2,$$

$$\frac{d\sigma_{FDM}}{dt} \propto \sigma_{FDM}^{h^*} \times \left| \frac{1}{t - m_{H_1}^2 + im_{H_1}\Gamma_{H_1}} - \frac{1}{t - m_{H_2}^2 + im_{H_2}\Gamma_{H_2}} \right|^2 \cdot (2t - 8m_\chi^2),$$

$$\frac{d\sigma_{VDM}}{dt} \propto \sigma_{VDM}^{h^*} \times \left| \frac{1}{t - m_{H_1}^2 + im_{H_1}\Gamma_{H_1}} - \frac{1}{t - m_{H_2}^2 + im_{H_2}\Gamma_{H_2}} \right|^2 \cdot \left(2 + \frac{(t - 2m_D^2)^2}{4m_V^4} \right).$$



Fix DM mass = 80 GeV, $\sin(\alpha) = 0.3$,
and vary H_2 mass (200,300,400,500) GeV

Asymptotic behavior in the full theory ($t \equiv m_{\chi\chi}^2$)

$$\text{ScalarDM : } G(t) \sim \frac{1}{(t - m_H^2)^2 + m_H^2 \Gamma_H^2} \quad (5.7)$$

$$\text{SFDM : } G(t) \sim \left| \frac{1}{t - m_1^2 + im_1 \Gamma_1} - \frac{1}{t - m_2^2 + im_2 \Gamma_2} \right|^2 (t - 4m_\chi^2) \quad (5.8)$$

$$\rightarrow \left| \frac{1}{t^2} \right|^2 \times t \sim \frac{1}{t^3} \quad (\text{as } t \rightarrow \infty) \quad (5.9)$$

$$\text{VDM : } G(t) \sim \left| \frac{1}{t - m_1^2 + im_1 \Gamma_1} - \frac{1}{t - m_2^2 + im_2 \Gamma_2} \right|^2 \left[2 + \frac{(t - 2m_V^2)^2}{4m_V^4} \right] \quad (5.10)$$

$$\rightarrow \left| \frac{1}{t^2} \right|^2 \times t^2 \sim \frac{1}{t^2} \quad (\text{as } t \rightarrow \infty) \quad (5.11)$$

Asymptotic behavior w/o the 2nd Higgs (EFT)

$$\text{SFDM : } G(t) \sim \frac{1}{(t - m_H^2)^2 + m_H^2 \Gamma_H^2} (t - 4m_\chi^2)$$

$$\rightarrow \frac{1}{t} \quad (\text{as } t \rightarrow \infty)$$



$$\text{VDM : } G(t) \sim \frac{1}{(t - m_H^2)^2 + m_H^2 \Gamma_H^2} \left[2 + \frac{(t - 2m_V^2)^2}{4m_V^4} \right]$$

$$\rightarrow \text{constant} \quad (\text{as } t \rightarrow \infty)$$

**Unitarity is
violated in EFT!**

Fermi-LAT GC γ -ray


see arXiv:1612.05687 for a recent overview by C.Karwin, S. Murgia, T. Tait, T.A.Porter,P.Tanedo




$$\text{GC} : b \sim l \lesssim 0.1^\circ$$

extended
GeV scale excess!

- **A DM interpretation**



DM + DM $\rightarrow b\bar{b}$ with $\sigma v = 1.7 \times 10^{-26} \text{cm}^3/\text{s}$



* See "1402.6703, T. Daylan et.al." for other possible channels

- **Millisecond Pulsars (astrophysical alternative)**


It may or may not be the main source, depending on

- luminosity func.
- bulge population
- distribution of bulge population

* See "1404.2318, Q. Yuan & B. Zhang" and "1407.5625, I. Cholis, D. Hooper & T. Linden"

GC gamma ray in HP VDM

P. Ko, WI Park, Y. Tang. arXiv:1404.5257, JCAP



H2 : 125 GeV Higgs
H1 : absent in EFT

Figure 2. Dominant s channel $b + \bar{b}$ (and $\tau + \bar{\tau}$) production




Figure 3. Dominant s/t -channel production of H_1 s that decay dominantly to $b + \bar{b}$

Importance of HP VDM with Dark Higgs Boson




Figure 4. Relic density of dark matter as function of m_ψ for $m_h = 125$, $m_\phi = 75$ GeV, $g_X = 0.2$, and $\alpha = 0.1$.




Figure 5. Illustration of γ spectra from different channels. The first two cases give almost the same spectra while in the third case γ is boosted so the spectrum is shifted to higher energy.

This mass range of VDM would have been impossible in the VDM model (EFT)

And No 2nd neutral scalar (Dark Higgs) in EFT

Summary

- Phenomenology of HP VDM and Singlet FDM presented within EFT vs. UV completed models
- EFT approach has a number of drawbacks : non-renormalizable, unitarity violation at high energy colliders, and it applies only if $m_{DM}, m_{SM} \ll m_\phi$ [But we don't know mass scales of dark particles !]
- In particular, one has $\Gamma_{\text{EFT}}(H_{125} \rightarrow VV) \rightarrow \infty$, as $m_V \rightarrow 0$, whereas it is finite in UV completed models [Importance of gauge invariance, unitarity and renormalizability]
- The dark Higgs ϕ can play crucial roles in interpreting the DM signatures at colliders, explaining the GC γ -ray excess ($VV \rightarrow \phi\phi$), improving vacuum stability up to Planck scale, modifying the Higgs inflation [ϕ should be actively searched for !]

Inelastic DM and XENON1T Excess

We consider Both Scalar and Fermion IDM



arXiv:2006.16876, PLB 810 (2020) 135848
With Seungwon Baek, Jongkuk Kim

XENON1T Excess

This excess disappeared recently, but we learn important lessons !

- Excess between 1-7 keV
 - Expected : 232 ± 15 , Observed : 285
 - Deviation $\sim 3.5 \sigma$
- Tritium contamination
 - Long half lifetime (12.3 years)
 - Abundant in atmosphere and cosmogenically produced in Xenon
- Solar axion
 - Produced in the Sun
 - Favored over bkgd @ 3.5σ
- Neutrino magnetic dipole moment
 - Favored @ 3.2σ



Electron recoil



DD/CMB Constraints

- To evade stringent bounds from direct detection expt's : sub GeV DM
- CMB bound excludes thermal DM freeze-out determined by S-wave annihilation : DM annihilation should be mainly in P-wave $\langle\sigma v\rangle \sim a + bv^2$

Planck 2018
R.K.Leane 35 al, PRD2018



Exothermic DM

- Inelastic exothermic scattering of XDM
- $XDM + e_{\text{atomic}} \rightarrow DM + e_{\text{free}}$ by dark photon exchange + kinetic mixing
- Excess is determined by $E_R \sim \delta = m_{XDM} - m_{DM}$
- Most works are based on effective/toy models where δ is put in by hand, or ignored dark Higgs
- dim-2 op for scalar DM and dim-3 op for fermion DM : soft and explicit breaking of local gauge symmetry, and include massive dark photon as well \rightarrow theoretically inconsistent !

Z_2 DM models with dark Higgs

- We solve this inconsistency and unitarity issue with Krauss-Wilczek mechanism
- By introducing a dark Higgs, we have many advantages:
 - Dark photon gets massive
 - Mass gap δ is generated by dark Higgs mechanism
 - We can have DM pair annihilation in P-wave involving dark Higgs in the final states, unlike in other works

Usual Approaches

For example, Harigaya, Nagai, Suzuki, arXiv:2006.11938

$$V(\phi) = m^2|\phi|^2 + \Delta^2 (\phi^2 + \phi^{*2}), \quad (1)$$

This term is problematic

$$\mathcal{L} = g_D A'^{\mu} (\chi_1 \partial_{\mu} \chi_2 - \chi_2 \partial_{\mu} \chi_1) + \epsilon e A'_{\mu} J_{\text{EM}}^{\mu},$$

Similarly for the fermion DM case





FIG. 1. Inelastic scattering of the heavier DM particle χ_2 off the electron e into the lighter particle χ_1 , mediated by the dark photon A' .

- The model is not mathematically consistent, since there is no conserved current a dark photon can couple to in the massless limit
- The second term with Δ^2 breaks $U(1)_X$ explicitly, although softly



For example, Harigaya, Nagai, Suzuki, arXiv:2006.11938




FIG. 4. The required value of ϵ to explain the observed excess of events at XENON1T in terms of the dark photon mass $m_{A'}$ (black solid lines). The left and right panels correspond to the cases of $m > m_{A'}/2$ and $m < m_{A'}/2$ respectively. We assume $g_D = 1.2$ in both cases. The blue lines denote the required value of ϵ to obtain the observed DM abundance by the thermal freeze-out process, discussed in Sec. IV. The solid lines correspond to the case without any entropy production. The dashed lines assume freeze-out during a matter dominated era and the subsequent reheating at T_{RH} , which suppresses the DM abundance by a factor of $(T_{RH}/T_{FO})^3$. The black dashed lines denote the mass density of χ_2 normalized by the total DM density. The shaded regions show the constraints from dark radiation and various searches for the dark photon A' which are discussed in Sec. V.

**Without dark Higgs, one can not consider light FDM,
because of CMB bounds on the S-wave annihilation**

Scalar XDM (X_R & X_I)

Field	ϕ	X	χ
U(1) charge	2	1	1

$$\begin{aligned}
 \mathcal{L} = & \mathcal{L}_{\text{SM}} - \frac{1}{4} \hat{X}_{\mu\nu} \hat{X}^{\mu\nu} - \frac{1}{2} \sin \epsilon \hat{X}_{\mu\nu} \hat{B}^{\mu\nu} + D^\mu \phi^\dagger D_\mu \phi + D^\mu X^\dagger D_\mu X - m_X^2 X^\dagger X + m_\phi^2 \phi^\dagger \phi \\
 & - \lambda_\phi (\phi^\dagger \phi)^2 - \lambda_X (X^\dagger X)^2 - \lambda_{\phi X} X^\dagger X \phi^\dagger \phi - \lambda_{\phi H} \phi^\dagger \phi H^\dagger H - \lambda_{HX} X^\dagger X H^\dagger H \\
 & - \mu (X^2 \phi^\dagger + H.c.), \tag{1}
 \end{aligned}$$

$$X = \frac{1}{\sqrt{2}}(X_R + iX_I),$$

$$H = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}(v_H + h_H) \end{pmatrix}, \quad \phi = \frac{1}{\sqrt{2}}(v_\phi + h_\phi),$$

$$\mathcal{L} \supset \epsilon g_X s_W Z^\mu (X_R \partial_\mu X_I - X_I \partial_\mu X_R) - \frac{g_Z}{2} Z_\mu \bar{\nu}_L \gamma^\mu \nu_L$$

$$\mathcal{L} \supset g_X Z'^\mu (X_R \partial_\mu X_I - X_I \partial_\mu X_R) - \epsilon e c_W Z'_\mu \bar{e} \gamma^\mu e,$$

$$U(1) \rightarrow Z_2 \text{ by } v_\phi \neq 0 : X \rightarrow -X$$




FIG. 1: (*left*) Feynman diagrams relevant for thermal relic density of DM: $XX^\dagger \rightarrow Z'\phi$ and (*right*) the region in the $(m_{Z'}, \epsilon)$ plane that is allowed for the XENON1T electron recoil excess and the correct thermal relic density for scalar DM case for $\delta = 2$ keV : (a) $m_{\text{DM}} = 0.1$ GeV. Different colors represents $m_\phi = 20, 40, 60, 80$ MeV. The gray areas are excluded by various experiments, from BaBar [61], E774 [62], E141 [63], Orasay [64], and E137 [65], assuming $Z' \rightarrow X_R X_I$ is kinematically forbidden.

P-wave annihilation x-sections

Scalar DM : $XX^\dagger \rightarrow Z'^* \rightarrow Z'\phi$

$$\sigma v \simeq \frac{g_X^4 v^2}{384\pi m_X^4 (4m_X^2 - m_{Z'}^2)^2} (16m_X^4 + m_{Z'}^4 + m_\phi^4 + 40m_X^2 m_{Z'}^2 - 8m_X^2 m_\phi^2 - 2m_{Z'}^2 m_\phi^2) \times \left[\{4m_X^2 - (m_{Z'} + m_\phi)^2\} \{4m_X^2 - (m_{Z'} - m_\phi)^2\} \right]^{1/2} + \mathcal{O}(v^4), \quad (10)$$

Fermion XDM (χ_R & χ_I)

$$\mathcal{L} = -\frac{1}{4}\hat{X}^{\mu\nu}\hat{X}_{\mu\nu} - \frac{1}{2}\sin\epsilon\hat{X}_{\mu\nu}B^{\mu\nu} + \bar{\chi}(i\not{D} - m_\chi)\chi + D_\mu\phi^\dagger D^\mu\phi - \mu^2\phi^\dagger\phi - \lambda_\phi|\phi|^4 - \frac{1}{\sqrt{2}}\left(y\phi^\dagger\bar{\chi}^c\chi + \text{h.c.}\right) - \lambda_{\phi H}\phi^\dagger\phi H^\dagger H$$

$$\begin{aligned}\chi &= \frac{1}{\sqrt{2}}(\chi_R + i\chi_I), \\ \chi^c &= \frac{1}{\sqrt{2}}(\chi_R - i\chi_I), \\ \chi_R^c &= \chi_R, \quad \chi_I^c = \chi_I,\end{aligned}$$

$$\begin{aligned}\mathcal{L} &= \frac{1}{2}\sum_{i=R,I}\bar{\chi}_i(i\not{D} - m_i)\chi_i - i\frac{g_X}{2}(Z'_\mu + \epsilon_{SW}Z_\mu)(\bar{\chi}_R\gamma^\mu\chi_I - \bar{\chi}_I\gamma^\mu\chi_R) \\ &\quad - \frac{1}{2}yh_\phi(\bar{\chi}_R\chi_R - \bar{\chi}_I\chi_I),\end{aligned}$$

$$U(1) \rightarrow Z_2 \text{ by } v_\phi \neq 0 : \chi \rightarrow -\chi$$





FIG. 2: (*top*) Feynman diagrams for $\chi\bar{\chi} \rightarrow \phi\phi$. (*bottom*) the region in the (m_Z, ϵ) plane that is allowed for the XENON1T electron recoil excess and the correct thermal relic density for fermion DM case for $\delta = 2$ keV and the fermion DM mass to be $m_R = 10$ MeV. Different colors represents $m_\phi = 2, 4, 6, 8$ MeV. The gray areas are excluded by various experiments, assuming $Z' \rightarrow \chi_R\chi_L$ is kinematically allowed, and the experimental constraint is weaker in the ϵ we are interested in, compared with the scalar DM case in Fig. 1 (right). We also show the current experimental bounds by NA64 [66].

P-wave annihilation x-sections

Scalar DM : $XX^\dagger \rightarrow Z'^* \rightarrow Z'\phi$

$$\sigma v \simeq \frac{g_X^4 v^2}{384\pi m_X^4 (4m_X^2 - m_{Z'}^2)^2} (16m_X^4 + m_{Z'}^4 + m_\phi^4 + 40m_X^2 m_{Z'}^2 - 8m_X^2 m_\phi^2 - 2m_{Z'}^2 m_\phi^2) \\ \times \left[\{4m_X^2 - (m_{Z'} + m_\phi)^2\} \{4m_X^2 - (m_{Z'} - m_\phi)^2\} \right]^{1/2} + \mathcal{O}(v^4), \quad (10)$$

Fermion DM : $\chi\bar{\chi} \rightarrow \phi\phi$

$$\sigma v = \frac{y^2 v^2 \sqrt{m_\chi^2 - m_\phi^2}}{96\pi m_\chi} \left[\frac{27\lambda_\phi^2 v_\phi^2}{(4m_\chi^2 - m_\phi^2)^2} + \frac{4y^2 m_\chi^2 (9m_\chi^4 - 8m_\chi^2 m_\phi^2 + 2m_\phi^4)}{(2m_\chi^2 - m_\phi^2)^4} \right] + \mathcal{O}(v^4), \quad (28)$$

Crucial to include “dark Higgs” to have Light DM pair annihilation in P-wave

Summary

- Local Z_2 scalar/fermion DM : theoretically well defined & mathematically consistent models for XDM
- Can explain a number of phenomena including the recent XENON1T data
- One can discriminate the spin of (X)DM at Belle2 from the polar angle distributions of the decaying points
- DM mass and the Δm can be determined with the focus point method
- Similar studies at ILC, CEPC, HL-LHC and FCC-hh in progress (The current version of FCC CDR does not include this interesting case.)

$U(1)_{L_\mu - L_\tau}$ -charged DM


: Z' only vs. $Z' + \phi$

arXiv:2204.04889 [hep-ph]
With Seungwon Baek, Jongkuk Kim

SM+ $U(1)_{L_\mu-L_\tau}$ gauge sym

- He, Josh, Lew, Volkas, PRD 43, 22; PRD 44, 2118 (1991)
- One of the anomaly free gauge groups without extension of fermion contents
- The simplest anomaly free U(1) extensions that couple to the SM fermions directly
- Can affect the muon g-2, PAMELA e^+ excess, (and B anomalies with extra fermions : Not covered in this talk)

Muon g-2



The Muon g-2 Collaboration, 2104.03281

Excellent example for graduate students

- Relativistic E&M (spinning particle in EM fields)
- Special relativity (time dilation)
- (V-A) structure of charged weak interaction

Muon (g-2) in $U(1)_{\mu-\tau}$ Model

Baek, Deshpande, He, Ko : hep-ph/0104141

Baek, Ko : arXiv:0811.1646 [hep-ph]

$$\begin{array}{ll} L_L^e : (1, 2, -1)(0) & e_R : (1, 1, -2)(0) \\ L_L^\mu : (1, 2, -1)(2a) & \mu_R : (1, 1, -2)(2a) \\ L_L^\tau : (1, 2, -1)(-2a) & \mu_R : (1, 1, -2)(-2a) \end{array}$$

$$\Delta a_\mu = \frac{\alpha'}{2\pi} \int_0^1 dx \frac{2m_\mu^2 x^2 (1-x)}{x^2 m_\mu^2 + (1-x)M_{Z'}^2} \approx \frac{\alpha'}{2\pi} \frac{2m_\mu^2}{3M_{Z'}^2}$$

$$Z' \rightarrow \mu^+ \mu^-, \tau^+ \tau^-, \nu_\alpha \bar{\nu}_\alpha \text{ (with } \alpha = \mu \text{ or } \tau), \psi_D \bar{\psi}_D$$

$\Gamma(Z' \rightarrow \mu^+ \mu^-) = \Gamma(Z' \rightarrow \tau^+ \tau^-) = 2\Gamma(Z' \rightarrow \nu_\mu \bar{\nu}_\mu) = 2\Gamma(Z' \rightarrow \nu_\tau \bar{\nu}_\tau) = \Gamma(Z' \rightarrow \psi_D \bar{\psi}_D)$
if $M_{Z'} \gg m_\mu, m_\tau, M_{DM}$. The total decay rate of Z' is approximately given by

$$\Gamma_{\text{tot}}(Z') = \frac{\alpha'}{3} M_{Z'} \times 4(3) \approx \frac{4(\text{or } 3)}{3} \text{ GeV} \left(\frac{\alpha'}{10^{-2}} \right) \left(\frac{M_{Z'}}{100\text{GeV}} \right)$$

$$\begin{array}{l} q\bar{q} \text{ (or } e^+e^-) \rightarrow \gamma^*, Z^* \rightarrow \mu^+ \mu^- Z', \tau^+ \tau^- Z' \\ \rightarrow Z^* \rightarrow \nu_\mu \bar{\nu}_\mu Z', \nu_\tau \bar{\nu}_\tau Z' \end{array}$$





FIG. 2. Δa_μ on the a vs. $m_{Z'}$ plane in case b). The lines from left to right are for Δa_μ away from its central value at $+2\sigma, +1\sigma, 0, -1\sigma$ and -2σ , respectively.

Baek and Ko, arXiv:0811.1646, for PAMELA e^+ excess

$$\mathcal{L}_{\text{Model}} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{New}}$$

$$\begin{aligned} \mathcal{L}_{\text{New}} = & -\frac{1}{4} Z'_{\mu\nu} Z'^{\mu\nu} + \bar{\psi}_D i D \cdot \gamma \psi_D - M_{\psi_D} \bar{\psi}_D \psi_D + D_\mu \phi^* D^\mu \phi \\ & - \lambda_\phi (\phi^* \phi)^2 - \mu_\phi^2 \phi^* \phi - \lambda_{H\phi} \phi^* \phi H^\dagger H. \end{aligned}$$


Here we ignored kinetic mixing for simplicity

$$D_\mu = \partial_\mu + ieQ A_\mu + i \frac{e}{s_W c_S} (I_3 - s_W^2 Q) Z_\mu + ig' Y' Z'_\mu$$

muon $g-2$, Leptophilic DM, Collider Signature

Muon ($g-2$)

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (302 \pm 88) \times 10^{-11}.$$



$$\Delta a_\mu = \frac{\alpha'}{2\pi} \int_0^1 dx \frac{2m_\mu^2 x^2 (1-x)}{x^2 m_\mu^2 + (1-x)M_{Z'}^2} \approx \frac{\alpha'}{2\pi} \frac{2m_\mu^2}{3M_{Z'}^2}$$




Figure 1: The relic density of CDM (black), the muon $(g-2)_\mu$ (blue band), the production cross section at B factories (1 fb, red dotted), Tevatron (10 fb, green dotdashed), LEP (10 fb, pink dotted), LEP2 (10 fb, orange dotted), LHC (1 fb, 10 fb, 100 fb, blue dashed) and the Z^0 decay width (2.5×10^{-6} GeV, brown dotted) in the $(\log_{10} \alpha', \log_{10} M_{Z'})$ plane. For the relic density, we show three contours with $\Omega h^2 = 0.106$ for $M_{\psi_D} = 10$ GeV, 100 GeV and 1000 GeV. The blue band is allowed by $\Delta a_\mu = (302 \pm 88) \times 10^{-11}$ within 3σ .




FIG. 1. The leading order contribution of the Z' to neutrino trident production (another diagram with μ^+ and μ^- reversed in g'




FIG. 2. Parameter space for the Z' gauge boson. The light-grey area is excluded at 95% C.L. by the CCFR measurement of the neutrino trident cross-section. The grey region with the dotted contour is excluded by measurements of the SM

Seungwon Baek, Pyungwon Ko,
arXiv:0811.1646, JCAP(2009)
about PAMELA e^+ excess

Altmannshofer et al.
arXiv:1406.2332 [hep-ph]

Neutrino trident puts strong
constraints on this model

One can evade the neutrino trident constraint, if one introduces
New fermions and generate muon $g-2$ at loop level w/ new fermions !

Z' Only

- Consider light Z' and $g_X \sim (\text{a few}) \times 10^{-4}$ for the muon g-2. Then
- $\chi\bar{\chi} \rightarrow Z'^* \rightarrow f_{\text{SM}}\bar{f}_{\text{SM}}$: dominant annihilation channel
- $g_X \sim 10^{-4}$ is too small for $\chi\bar{\chi} \rightarrow Z'Z'$ to be effective for $\Omega_\chi h^2$
- $m_{Z'} \sim 2m_{\text{DM}}$ with the s-channel Z' resonance for the correct relic density
- Many recent studies on this case:
 - Asai, Okawa, Tsumura, 2011.03165
 - Holst, Hooper, Krnjaic, 2107.09067
 - Drees and Zhao, arXiv:2107.14528
 - And some earlier papers




FIG. 1. Regions inside the yellow and Green shaded areas by the Δa_μ are allowed at 1σ and 2σ C.L.. Cyan, black, and orange regions are excluded by other experimental bounds. Above green solid line is ruled out by the Borexino experiment. Region inside the orange area can resolve the Hubble tension. We take two Benchmark Points (BP) $(M_{Z'}, g_X)$ as **BPI** = $(11.5 \text{ MeV}, 4 \times 10^{-4})$ and **BPII** = $(100 \text{ MeV}, 8 \times 10^{-4})$.

$U(1)_{L_\mu - L_\tau}$ -charged DM

: Z' only vs. $Z' + \phi$

cf: Let me call Z' , $U(1)_{L_\mu - L_\tau}$ gauge boson,
“dark photon”, since it couples to DM

Models with Φ

TABLE I: $U(1)$ charge assignments of newly introduced particles and SM particles. The other SM particles are singlet.

Field	Z'_μ	$X(\chi)$	Φ	$L_\mu = (\nu_{L\mu}, \mu_L), \mu_R$	$L_\tau = (\nu_{L\tau}, \tau_L), \tau_R$
spin	1	0 (1/2)	0	1/2	1/2
$U(1)$ charge	0	$Q_X(Q_\chi)$	Q_Φ	+1	-1

We Consider Both Complex Scalar (X) and Dirac Fermion DM (χ)

- Physics depends on Q_Φ , Q_X and Q_χ
- $Q_\Phi = 2Q_{X(\chi)}$ and $3Q_X$ need special cares, since there are extra gauge invariant op's that break $U(1) \rightarrow Z_2, Z_3$ after $U(1)$ is spontaneously broken by nonzero VEV of Φ

Complex Scalar DM (generic with $Q_\Phi \neq Q_X$, etc)




FIG. 2. (Top) Feynman diagrams for Complex scalar DM annihilating to a pair of Z' bosons. (Bottom) Feynman diagrams for Complex scalar DM annihilating to a pair of H_1 bosons.

$$H_2 \simeq H_{125} \text{ and } H_1 \simeq \phi \text{ (dark Higgs)}$$




FIG. 3. *Top*: relic abundance of complex scalar DM as functions of $\lambda_{\phi X}$ for [BPI] for $M_X = 1, 10, 100, 1000$ GeV, respectively. We assumed $Q_\Phi = 1.1$, $M_{H_1} = 1$ GeV, and $\sin \alpha = 10^{-4}$. Solid (Dashed) lines represent the region where bounds on DM direct detection are satisfied (ruled out). *Bottom*: the preferred parameter space in the $(M_{H_1}, \lambda_{\phi X})$ plane for $\lambda_{HX} = 0$.




FIG. 7. The (*Top*) plots show the relic abundance of complex scalar DM for $Q_\Phi = 1.1$ as functions of dark Higgs mass M_{H_1} for [BPI] (*Left*) and [BPII] (*Right*). The (*Bottom*) plots show the relic density as functions of $\lambda_{\phi X}$ (*Left*) and the preferred parameter space in the $(M_{H_1}, \lambda_{\phi X})$ plane for $\lambda_{HX} = 0$ (*Right*) for [BPII]. We take four different DM masses, $M_X = 1, 10, 100, 1000$ GeV, respectively. Solid (Dashed) lines represent the region where bounds on DM direct detection are satisfied (ruled out).

DM mass : much wider range than $m_{Z'} \sim 2m_{\text{DM}}$
due to dark Higgs boson contributions

Complex Scalar DM:

$$U(1)_{L_\mu - L_\tau} \rightarrow Z_2 \quad (Q_\Phi = 2Q_X)$$




FIG. 8. (Top) Feynman diagrams for local Z_2 scalar DM annihilating to a pair of Z' bosons. (Bottom) Feynman diagrams for local Z_2 scalar DM annihilating to a pair of H_1 bosons, which is mostly dark Higgs-like.




FIG. 4. *Top*: Relic abundance of local Z_2 scalar DM as functions of $\lambda_{\Phi X}$ for [BPI] and different values of mass splittings (Δ). We take $\lambda_{HX} = 0$, $M_{H_1} = 10\text{GeV}$, and $s_\alpha = 10^{-4}$. All the curves satisfy the DM direct detection bound. *Bottom*: The preferred parameter space in the $(M_{H_1}, \lambda_{\Phi X})$ plane for different values of Δ . The gray area is excluded by the perturbative condition.




FIG. 9. (*Left*) Relic abundance of local Z_2 scalar DM in case of [BPII]. We take $\lambda_{HX} = 0$, $M_{H_1} = 10\text{GeV}$, and $s_\alpha = 10^{-4}$. All the lines satisfy the DM direct detection bound. (*Right*) Relic abundance of local Z_2 scalar DM in the $(M_{H_1}, \lambda_{\Phi X})$ plane.

**DM mass : much wider range than $m_{Z'} \sim 2m_{\text{DM}}$
due to dark Higgs boson contributions**

Complex Scalar DM:

$$U(1)_{L_\mu - L_\tau} \rightarrow Z_3 \quad (Q_\Phi = 3Q_X)$$

**Local Z_3 DM Model : first considered by Ko, Tang:
arXiv:1402.6449 (SIDM), 1407.5492 (GC γ -ray excess)**

$$V = -\mu_H^2 H^\dagger H + \lambda_H (H^\dagger H)^2 - \mu_\phi^2 \phi_X^\dagger \phi_X + \lambda_\phi (\phi_X^\dagger \phi_X)^2 + \mu_X^2 X^\dagger X + \lambda_X (X^\dagger X)^2 \\ + \lambda_{\phi H} \phi_X^\dagger \phi_X H^\dagger H + \lambda_{\phi X} X^\dagger X \phi_X^\dagger \phi_X + \lambda_{HX} X^\dagger X H^\dagger H + (\lambda_3 X^3 \phi_X^\dagger + H.c.) \quad (2.1)$$





FIG. 1: Feynman diagrams for dark matter semi-annihilation. Only (a), (b), and (c) with H_1 as final state appear in the global Z_3 model, while all diagrams could contribute in local Z_3 model.

**ϕ and Z' : present only in models with dark gauge symmetries,
And not in models with global dark symmetries**

Dark $U(1) \rightarrow Z_3$

$\Omega h^2 \subset [0.1145, 0.1253], \lambda_3 < 0.02$



- Blue band marks the upper bound,
- All points are allowed in our local Z_3 model, 1402.6449
- only circles are allowed in global Z_3 model, 1211.1014

This whole region is allowed in local Z_3 case

$$r \equiv \frac{v\sigma^{XX \rightarrow X^*Y}}{2v\sigma^{XX^* \rightarrow YY} + \frac{1}{2}v\sigma^{XX \rightarrow X^*Y}}$$

$$U(1)_{L_\mu - L_\tau} \rightarrow Z_3$$




FIG. 10. Relic abundance of Z_3 scalar DM for the [BPI] (Left) and the [BPII] (Right), respectively. Here we fixed $\lambda_{HX} = 0$ for simplicity.

- $g_X \sim O(10^{-4})$: very small. $XX \rightarrow X^\dagger Z'$ is not important
DM mass : much wider range than $m_{Z'} \sim 2m_{\text{DM}}$
due to dark Higgs boson contributions
- λ_3 controlling $XX \rightarrow X^\dagger H_1$ is an important parameter

Dirac fermion DM:

$$U(1)_{L_\mu - L_\tau} \rightarrow Z_2 \quad (Q_\Phi = 2Q_\chi)$$




FIG. 5. Feynman diagrams of local Z_2 fermion DM (co-)annihilating into a pair of Z' bosons and H_1 bosons (*Top*), and $Z' + H_1$ (*Bottom*).




FIG. 6. *Top*: Dark matter relic density as functions of mass splitting Δ for [BPI] and for different values of DM mass, $M_I = 1, 10, 100, 1000 \text{ GeV}$. Solid (Dashed) lines denote the region where bounds on DM direct detection are satisfied (ruled out). *Bottom*: Preferred parameter space in the (M_{H_1}, Δ) plane for different DM masses. The gray region is ruled out by the perturbativity condition on λ_Φ .




FIG. 11. (*Top*) Dark matter relic density as functions of dark Higgs mass M_{H_1} for [BPI] (*Left*) and [BPII] (*Right*) (*Bottom-Left*) Dark matter relic density as functions of Δ for [BPII], and (*Bottom-right*) Preferred parameter region in the (Δ, M_{H_1}) plane. Solid (Dashed) lines denote the region where bounds on DM direct detection are satisfied (ruled out).

DM mass : much wider range than $m_{Z'} \sim 2m_{\text{DM}}$ due to dark Higgs boson contributions

Conclusion

- DM physics with massive dark photon can not be complete without including dark gauge symmetry breaking mechanism, e.g. dark Higgs field ϕ , which have been largely ignored by DM community (or some ways other than dark Higgs to provide dark photon mass)
- Many examples show the importance of ϕ in DM phenomenology, astroparticle physics and cosmology
- Once ϕ is included, can accommodate the muon $g-2$ and thermal DM without the s-channel resonance condition $m_{Z'} \sim 2m_{\text{DM}}$
- m_{DM} : essentially free, whereas $m_{Z'} \sim O(10 - 100)$ MeV and $g_X \sim O(10^{-4})$ can explain the muon ($g-2$)

Higgs-Portal Assisted Higgs Inflation

**arXiv:1405.1635 [hep-ph], JCAP02 (2017) 003
With Jinsu Kim, Wan-Il Park**

Higgs Inflation

- Inflation : the main paradigm for very early Universe
- But no very compelling inflation scenarios based on high energy physics
- SM Higgs boson can play a role of inflaton if it has large non minimal coupling [Bezrukov, Shaposhnikov (2007)] or non canonical kinetic term
- Merits: Minimal model, Consistent with Planck data, Can connect low energy (m_{EW}) scale to high energy (inflation) scale

Higgs Inflation in SM

(before BICEP2)

[Bezrukov and Shaposhnikov, 0710.3755]

$$\frac{\mathcal{L}}{\sqrt{-g}} = -\frac{1}{2\kappa} \left(1 - \xi \frac{h^2}{M_{\text{Pl}}^2} \right) R + \mathcal{L}_h$$

Nonminimal coupling

$$\epsilon = \frac{M_{\text{P}}^2}{2} \left(\frac{dU/d\chi}{U} \right)^2 \simeq \frac{4M_{\text{P}}^4}{3\xi^2 h^4}$$

$$U(\chi) = \frac{1}{\Omega(\chi)^4} \frac{\lambda}{4} (h(\chi)^2 - v^2)^2$$

$$\eta = M_{\text{P}}^2 \frac{d^2 U/d\chi^2}{U} \simeq -\frac{4M_{\text{P}}^2}{3\xi h^2},$$


$$\Rightarrow \epsilon \simeq \frac{3}{4} \eta^2$$

$$n_s = 1 - 6\epsilon + 2\eta \sim 0.96$$

$$\Rightarrow \eta \simeq \frac{1}{2} (n_s - 1)$$

$$\Rightarrow \epsilon \simeq \frac{3}{16} (n_s - 1)^2$$

$$\Rightarrow r \simeq 16\epsilon \simeq 3 (n_s - 1)^2 \sim 5 \times 10^{-3}$$




Higgs Inflation in SM


(after BICEP2)

$r_{\text{BICEP2}} \sim 0.1$ \Rightarrow Is Higgs inflation ruled out? **No!**

$$U(h) = \frac{\lambda}{4\Omega^4} (h^2 - v_H^2)^2 \rightarrow \frac{\lambda(\mu)}{4\Omega^4} (h^2 - v_H^2)^2$$

[Hamada, Kawai, Oda and Park, I403.5043; Bezrukov and Shposhnikov, I403.6078]





Effects of running on slow-roll parameters



$$\epsilon = \frac{M_{\text{Pl}}^2}{2} \left(\frac{dh}{d\chi} \frac{dU}{dh} \right)^2 = \frac{1}{2} \left(4 + \frac{\beta_\lambda}{\lambda_H} \right)^2 \frac{M_{\text{Pl}}^2/h^2}{\sqrt{\Omega^2 + 6\xi^2 h^2/M_{\text{Pl}}^2}} \approx \frac{1}{12} \left(4 + \frac{\beta_\lambda}{\lambda_H} \right)^2 \frac{M_{\text{Pl}}^4}{\xi^2 h^4} \quad (17)$$

$$\begin{aligned} \eta &= \frac{M_{\text{Pl}}^2}{U} \frac{dh}{d\chi} \frac{d}{dh} \left(\frac{dh}{d\chi} \frac{dU}{dh} \right) \\ &= \left(4 + \frac{\beta_\lambda}{\lambda_H} \right) \frac{M_{\text{Pl}}^2}{h^2} \frac{\Omega^2}{\Omega^2 + 6\xi^2 h^2/M_{\text{Pl}}^2} \left\{ \frac{1}{\Omega^2} \frac{\beta_\lambda}{\lambda_H} \left[1 + \frac{d \ln(\beta_\lambda/\lambda_H)/d \ln \varphi}{4 + \beta_\lambda/\lambda_H} \right] + 3 - 2 \frac{d \ln \Omega^2}{d \ln h} - \frac{\xi(1+6\xi)h^2/M_{\text{Pl}}^2}{1 + \xi(1+6\xi)h^2/M_{\text{Pl}}^2} \right\} \\ &\simeq -\frac{1}{3} \left(4 + \frac{\beta_\lambda}{\lambda_H} \right) \frac{M_{\text{Pl}}^2}{\xi h^2} \left\{ 1 - \frac{M_{\text{Pl}}^2}{2\xi h^2} \frac{\beta_\lambda}{\lambda_H} \left[1 + \frac{d \ln \beta_\lambda/d \ln \varphi - \beta_\lambda/\lambda_H}{4 + \beta_\lambda/\lambda_H} \right] \right\} \end{aligned} \quad (18)$$

ϵ & η are independent



However m_t and m_h are tightly constrained!

CMS Preliminary



[CMS PAS HIG-13-001]

[1406.3827]



In SM, our vacuum is likely to be meta-stable.

Higgs portal interaction

$$V \supset \lambda_{\Phi H} |\Phi|^2 H^\dagger H \quad \longrightarrow \quad \lambda_H = \left[1 - \left(1 - \frac{m_\phi^2}{m_h^2} \right) \sin^2 \alpha \right] \lambda_H^{\text{SM}}$$

Scalar mixing



→ $\lambda_H > \lambda_H^{\text{SM}}$ for $m_\phi > m_h$ & $\alpha \neq 0$

→ Vacuum instability along the Higgs direction is easily removed.

→ Higgs inflation consistent with BICEP2 is possible for a wide range of m_t and M_h


Higgs portal interaction disconnect m_t and M_h from inflationary observables.

Higgs-portal Higgs inflation



$m_t = 173.2 \text{ GeV}$
 $M_h = 125.5 \text{ GeV}$

* Inflection point control
 $(\alpha, m_\phi) \ \& \ \lambda_{\Phi H}$



Result of numerical analysis

$k_* \times \text{Mpc}$	N_e	h_*/M_{Pl}	ϵ_*	η_*	$10^9 P_S$	n_s	r
0.002	59	0.83	0.00448	-0.02465	2.2639	0.9238	0.0717
0.05	56	0.72	0.00525	-0.00190	2.1777	0.9647	0.0840

- Result depends very sensitively on α , m_ϕ and $\lambda_{\Phi H}$ -

**Higgs Portal Higgs Inflation
 can have $r \lesssim O(0.1)$ without
 resorting to m_t and M_h .**


Singlet fermion CDM

Baek, Ko, Park, arXiv:1112.1847

$$\mathcal{L} = \mathcal{L}_{\text{SM}} - \mu_{HS} S H^\dagger H - \frac{\lambda_{HS}}{2} S^2 H^\dagger H + \frac{1}{2} (\partial_\mu S \partial^\mu S - m_S^2 S^2) - \mu'_S S - \frac{\mu''_S}{3} S^3 - \frac{\lambda_S}{4} S^4 + \bar{\psi} (i \not{\partial} - m_{\psi_0}) \psi - \lambda S \bar{\psi} \psi$$

mixing

invisible decay



Production and decay rates are suppressed relative to SM.

⦿ This simple model has not been studied properly !!

HP assisted HI w/ SFDM

$$\mathcal{L} \supset \frac{1}{2} \xi_s S^2 R$$

ξ_s term is generated by RG, even if $\xi_s = 0$ at $\mu = m_t$ scale.
We assume $S = 0$ during inflation : Inflation along the Higgs direction.




Figure 3. Jordan-frame Higgs potential V_{eff} (left panel) and the running of λ_H (right panel) in SFDM for $\xi_h = 440$, $\xi_s = 0$, $m_s = 600$ GeV, $\lambda_{SH} = 0.1$, $\lambda_S = 0.2$, and $\lambda_\psi = 0.3$ chosen at M_t scale.

HP assisted HI w/ SFDM

α	m_s	λ_{SH}	λ_S	λ_ψ	ξ_h	N_e	$10^9 P_S$	n_s	r	α_s
0.036	500	0.1	0.2	0.3	433	57.3	2.2	0.9758	0.0926	-0.0003
0.03885	500	0.1	0.1	0.1	396	57.3	2.2	0.9775	0.0878	-0.0003




Figure 4. Tensor-to-scalar ratio as a function of the mixing angle α for $m_s = 300$ GeV, 400 GeV, 500 GeV and 600 GeV at the pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$. Here $\xi_s = 0$, $\lambda_{SH} = 0.1$, $\lambda_S = 0.2$, and $\lambda_\psi = 0.3$ at M_t scale are used. The nonminimal coupling of the SM Higgs to gravity, ξ_h , is chosen in such a way that the Planck normalization (3.8) is satisfied. The grey-dotted lines indicate the parameter region where the spectral index n_s becomes larger than 2σ Planck bound, $n_s \gtrsim 0.98$. Similar behaviors are found for different sets of model parameters.




Figure 6. Tensor-to-scalar ratio as a function of the mixing angle α for $m_s = 300$ GeV, 400 GeV, 500 GeV and 600 GeV, with the constraints discussed in the main text. The stringent upper bounds for a given m_s comes from the DM physics. The values of the other parameters are the same as in figure 4. Color-shaded regions (following the scheme of colored lines) are the excluded regions from the latest LUX experiment, corresponding to different dark Higgs masses.

**Large $r \sim O(0.1)$ is possible in HP assisted HI,
without tight connection to m_t, m_h**

Summary

Message to take home

- DM interacting with massive dark photon is a typical scenario in DM physics
- Very often, dark Higgs boson (or some mechanism to generate dark photon mass) has been ignored
- However, there are many examples that show importance of dark Higgs boson
- In this talk, I discussed the following examples:

- HP VDM: $\Gamma_{\text{inv}}(H \rightarrow VV)$ and GC γ -ray excess
- XENON1T excess in terms of exothermic scattering of inelastic DM (both scalar and fermion DM)
- $U(1)_{L_\mu - L_\tau}$ -charged scalar/fermion DM outside the $m_{Z'} \sim 2m_{\text{DM}}$ window
- Higgs-portal assisted Higgs inflation with large $r \sim O(0.1)$
- Additional dark Higgs (singlet-like scalar) : generic (in DM models with dark gauge symmetry), improves EW vac stability \gg should be actively search for @ LHC and other future colliders

BACKUP SLIDES


DM is stable because...

- Symmetries

- (ad hoc) Z_2 symmetry, etc
- R-parity
- Topology (from a broken sym.)

- Very small mass and weak coupling

e.g: QCD-axion ($m_a \sim \Lambda_{\text{QCD}}^2/f_a$; $f_a \sim 10^9\text{-}12$ GeV)


$$\Gamma_a \sim \mathcal{O}(10^{-5}) \frac{m_a^3}{f_a^2} \ll H_0 \sim 10^{-42} \text{ GeV}$$

But for WIMP ...

S.Baek, P.Ko, W.I.Park, arXiv:1303.4280 [hep-ph], JHEP (2013)

- Global symmetry is not enough since

$$-\mathcal{L}_{\text{int}} = \begin{cases} \lambda \frac{\phi}{M_{\text{P}}} F_{\mu\nu} F^{\mu\nu} & \text{for boson} \\ \lambda \frac{1}{M_{\text{P}}} \bar{\psi} \gamma^\mu D_\mu \ell_{Li} H^\dagger & \text{for fermion} \end{cases}$$

Observation requires [M.Ackermann et al. (LAT Collaboration), PRD 86, 022002 (2012)]

$$\tau_{\text{DM}} \gtrsim 10^{26-30} \text{sec} \Rightarrow \begin{cases} m_\phi \lesssim \mathcal{O}(10) \text{keV} \\ m_\psi \lesssim \mathcal{O}(1) \text{GeV} \end{cases}$$

\Rightarrow WIMP is unlikely to be stable

- SM is guided by gauge principle

It looks natural and may need to consider a gauge symmetry in dark sector, too.

Why Dark Gauge Symmetry ?

S.Baek, P.Ko, W.I.Park, arXiv:1303.4280 [hep-ph], JHEP (2013)

- Is DM absolutely stable or very long lived ?
- If DM is absolutely stable, one can assume it carries a new **conserved dark charge**, associated with **unbroken dark gauge sym**
- DM can be long lived (lower bound on DM lifetime is much weaker than that on proton lifetime) if dark sym is spontaneously broken

Higgs can be harmful to weak scale DM stability

Z₂ sym Scalar DM

$$\mathcal{L} = \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} m_S^2 S^2 - \frac{\lambda_S}{4!} S^4 - \frac{\lambda_{SH}}{2} S^2 H^\dagger H.$$

- Very popular alternative to SUSY LSP
- Simplest in terms of the # of new dof's
- But, where does this Z₂ symmetry come from ?
- Is it Global or Local ?

Fate of CDM with Z_2 sym

- Global Z_2 cannot save DM from decay with long enough lifetime

Consider Z_2 breaking operators such as

$$\frac{1}{M_{\text{Planck}}} SO_{\text{SM}}$$

keeping dim-4 SM operators only

The lifetime of the Z_2 symmetric scalar CDM S is roughly given by

$$\Gamma(S) \sim \frac{m_S^3}{M_{\text{Planck}}^2} \sim \left(\frac{m_S}{100\text{GeV}}\right)^3 10^{-37} \text{GeV}$$

The lifetime is too short for ~ 100 GeV DM


Fate of CDM with Z_2 sym

- Spontaneously broken local $U(1)_X$ can do the job to some extent, but there is still a problem

Let us assume a local $U(1)_X$ is spontaneously broken by $\langle \phi_X \rangle \neq 0$ with

$$Q_X(\phi_X) = Q_X(X) = 1$$

Then, there are two types of dangerous operators:



- These arguments will apply to all the CDM models based on ad hoc Z_2 symmetry
- One way out is to implement Z_2 symmetry as local $U(1)$ symmetry (arXiv:1407.6588 with Seungwon Baek and Wan-Il Park)
- See a paper by Ko and Tang on local Z_3 scalar DM, and another by Ko, Omura and Yu on inert 2HDM with local $U(1)_H$

$$Q_X(\phi) = 2, \quad Q_X(X) = 1$$

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_{\text{SM}} + -\frac{1}{4}X_{\mu\nu}X^{\mu\nu} - \frac{1}{2}\epsilon X_{\mu\nu}B^{\mu\nu} + D_\mu\phi_X^\dagger D^\mu\phi_X - \frac{\lambda_X}{4}\left(\phi_X^\dagger\phi_X - v_\phi^2\right)^2 + D_\mu X^\dagger D^\mu X - m_X^2 X^\dagger X \\ & - \frac{\lambda_X}{4}(X^\dagger X)^2 - (\mu X^2\phi^\dagger + H.c.) - \frac{\lambda_{XH}}{4}X^\dagger X H^\dagger H - \frac{\lambda_{\phi_X H}}{4}\phi_X^\dagger\phi_X H^\dagger H - \frac{\lambda_{XH}}{4}X^\dagger X\phi_X^\dagger\phi_X \end{aligned}$$

The lagrangian is invariant under $X \rightarrow -X$ even after $U(1)_X$ symmetry breaking.

Unbroken Local Z2 symmetry

$X_R \rightarrow X_I\gamma_h^*$ followed by $\gamma_h^* \rightarrow \gamma \rightarrow e^+e^-$ etc.

The heavier state decays into the lighter state

The local Z2 model is not that simple as the usual Z2 scalar DM model (also for the fermion CDM)